arXiv:1501.02149v2 [gr-qc] 7 Mar 2015

Cosmological perturbations in mimetic matter model

Jiro Matsumoto[1]*, Sergei D. Odintsov[2,3], and Sergey V. Sushkov[1]

[1]*Institute of Physics,*
Kazan Federal University,
Kremlevskaya Street 18,
Kazan 420008, Russia
[2]*Consejo Superior de Investigaciones Científicas,*
ICE/CSIC-IEEC, Campus UAB,
Facultat de Ciències,
Torre C5-Parell-2a pl,
E-08193 Bellaterra (Barcelona), Spain
[3]*Institució Catalana de Recerca i Estudis Avançats (ICREA),*
Barcelona, Spain

Abstract

We investigate the cosmological evolution of mimetic matter model with arbitrary scalar potential. The cosmological reconstruction , which is the way to construct a model for arbitrary evolutions of the scale factor, is explicitly done for different choices of potential. The cases that mimetic matter model shows the evolution as Cold Dark Matter(CDM), wCDM model, dark matter and dark energy with dynamical $Om(z)$, where $Om(z) \equiv [(H(z)/H_0)^2 - 1]/[(1+z)^3 - 1]$, or phantom dark energy with phantom-non-phantom crossing are presented in detail. The cosmological perturbations for such evolution are studied in mimetic matter model. For instance, the evolution behavior of the matter density contrast which is different from usual one, i.e. $\ddot{\delta} + 2H\dot{\delta} - \kappa^2\rho\delta/2 = 0$ is investigated. The possibility of peculiar evolution of δ in the model under consideration is shown. Special attention is paid to the behavior of matter density contrast near to future singularity where decay of perturbations may occur much earlier the singularity.

PACS numbers:

---

* E-mail address: jmatsumoto@kpfu.ru

I. INTRODUCTION

The existence of dark energy and dark matter is indirectly shown by the observations of Cosmic Microwave Background (CMB), Large Scale Structure (LSS) of the Universe, Super Novae of Type Ia (SNIa), and related observational probes. Dark energy is usually regarded as a cosmological constant Λ in the standard model of cosmology, ΛCDM model. However, in general it represents some energy which have a negative pressure with equation of state parameter being less than $-1/3$. On the other hand, dark matter, in particular Cold Dark Matter (CDM) means the nonrelativistic unknown matter. There are many candidates for dark energy (for review, see[1, 2]) and dark matter[3, 4]. However, it is not yet clear which model is viable one.

In this paper, we will consider mimetic matter model [5], which was first proposed as a model of dark matter. However, it was realized later that it can be treated as a model of dark energy by adding the potential term of the scalar field [6]. Mimetic matter model is a conformally invariant theory by making the physical metric as a product of an auxiliary metric and the contraction of an auxiliary metric and the kinetic term of the scalar field. Whereas, the equivalence between mimetic matter model and a scalar field model with a Lagrange multiplier [7] is shown in [8, 9]. Therefore, we treat mimetic matter model as a scalar field model with a Lagrange multiplier in this paper.

The contents of the paper are the following. In Sec. II, we consider the background evolution of the Universe in general mimetic matter model. The cosmological reconstruction of the model , which is the way to construct the action which may realize the given evolution history of the universe by the choice of potential, is done and ΛCDM-like era is explicitly executed for corresponding scalar potential. The evolution of the matter density perturbation is discussed in Sec. III. First, mimetic matter model without potential is used to evaluate whether mimetic dark matter also behaves as dark matter at the cosmological perturbations level. Next, we investigate the matter density perturbation in general mimetic matter model with scalar potential. In Sec. IV, the possibility of the phantom – non-phantom transition is explicitly demonstrated in mimetic matter model. The behavior of the cosmological perturbations is also discussed, especially the behavior close to finite-time future singularity. Concluding remarks are given in Sec. V. The units of $k_{\mathrm{B}} = c = \hbar = 1$ are used and gravitational constant $8\pi G$ is denoted by $\kappa^2 \equiv 8\pi/{M_{\mathrm{Pl}}}^2$ with the Planck mass of $M_{\rm Pl} = G^{-1/2} = 1.2 \times 10^{19}$GeV in this paper.

II. THE ACCELERATING EVOLUTION OF THE ISOTROPIC AND HOMOGENEOUS UNIVERSE

We consider the action of mimetic matter model[6]:

$$S = \int d^4x\sqrt{-g}\left[\frac{1}{2\kappa^2}R - V(\phi) + \lambda(g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi + 1)\right] + S_{\rm matter}, \tag{2.1}$$

where V is an arbitrary function of the scalar field ϕ and λ is a Lagrange multiplier. The Einstein equation obtained from Eq. (2.1) is

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = -\kappa^2 g_{\mu\nu}V(\phi) - 2\kappa^2\lambda\partial_\mu\phi\partial_\nu\phi + \kappa^2 g_{\mu\nu}\lambda(g^{\rho\sigma}\partial_\rho\phi\partial_\sigma\phi + 1) + \kappa^2 T_{\mu\nu}, \tag{2.2}$$

where $R_{\mu\nu} = \partial_\sigma\Gamma^\sigma_{\mu\nu} - \partial_\mu\Gamma^\sigma_{\nu\sigma} + \Gamma^\sigma_{\mu\nu}\Gamma^\rho_{\sigma\rho} - \Gamma^\sigma_{\mu\rho}\Gamma^\rho_{\nu\sigma}$ and $T_{\mu\nu}$ is the energy momentum tensor of the usual matter. On the other hand, the equation given by the variation with respect to the scalar field ϕ is the following one:

$$- V_{,\phi} - 2\nabla^\mu(\lambda\partial_\mu\phi) = 0, \tag{2.3}$$

where $V_{,\phi} \equiv dV(\phi)/d\phi$. The constraint equation given by the variation of λ is

$$g^{\mu\nu}\partial_\mu\phi\partial_\nu\phi + 1 = 0. \tag{2.4}$$

When the spatially flat Friedmann-Lemaitre-Robertson-Walker (FLRW) metric, $ds^2 = -dt^2 + a^2(t)\sum_{i=1}^3 dx^i dx^i)$, is taken, FLRW equations are written by

$$3H^2 = \kappa^2\rho - 2\kappa^2\lambda\dot\phi^2 + \kappa^2 V + \kappa^2\lambda(\dot\phi^2 - 1), \tag{2.5}$$

$$-2\dot H - 3H^2 = \kappa^2 w\rho - \kappa^2 V - \kappa^2\lambda(\dot\phi^2 - 1), \tag{2.6}$$

where $H \equiv \dot a(t)/a(t)$. ρ is the energy-density of the matter and w is the equation of state (EoS) parameter expressed by $w = p/\rho$. From Eqs. (2.3) and (2.4) we have

$$2\dot\lambda\dot\phi + 2\lambda\ddot\phi + 6H\lambda\dot\phi - V_{,\phi} = 0, \tag{2.7}$$

$$\dot\phi^2 = 1. \tag{2.8}$$

The equation of continuity of the matter is

$$\dot\rho + 3(1+w)H\rho = 0. \tag{2.9}$$

If w is constant, one can integrate Eq. (2.9) as

$$\rho = \rho_0 \left(\frac{a}{a_0}\right)^{-3(1+w)}, \tag{2.10}$$

where $\rho_0 a_0^{3(1+w)}$ is an integration constant. While, combining Eqs. (2.5), (2.6), and (2.8) it gives the equations

$$\lambda = \frac{1}{\kappa^2}\dot{H} + \frac{1}{2}(1+w)\rho \tag{2.11}$$

and

$$V = w\rho + \frac{1}{\kappa^2}(2\dot{H} + 3H^2). \tag{2.12}$$

Therefore, all the solutions for the variables will be found if Eq. (2.12) can be solved with respect to $a(t)$. From the other side, one can say that this model is able to realize arbitrary evolution history of the Universe by using the arbitrariness of the function $V(\phi)$. Indeed, one can choose the form of $V(\phi)$ for the arbitrary function f such that $a(t) = f(t)$ as

$$V(\phi) = w\rho_0 \left(\frac{f(\phi)}{f(\phi_0)}\right)^{-3(1+w)} + \frac{1}{\kappa^2}\left(2\frac{f''(\phi)}{f(\phi)} + \frac{f'(\phi)^2}{f(\phi)^2}\right). \tag{2.13}$$

A. Cosmological reconstruction

Some forms of the function $V(\phi)$ are considered in ref.[6]. Here, we consider some other examples of $V(\phi)$, which are adequate to the cosmological observations and yield the accelerating universe. In particularly, the form of $V(\phi)$ which can yield the phantom – non-phantom transition is treated in Sec. IV. In the following, we assume the value of EoS parameter of the usual matter as $p_{\mathrm{matter}}/\rho_{\mathrm{matter}} = w_{\mathrm{matter}} = 0$.

1. Mimetic wCDM model

wCDM model consists of cold dark matter and a fluid with EoS parameter $w < 0$. For the case $w = -1$, wCDM model is equivalent to the ΛCDM model, so its background evolution can be realized by choosing the form of the potential as $V = \Lambda$. In the case of $w \neq -1$, we can reproduce the background evolution of the Universe in wCDM model if the effective energy density $-2\lambda + V$ and the effective pressure $-V$ satisfy the following conditions:

$$-2\lambda + V = \rho_{dm0} a^{-3} + \rho_{w0} a^{-3(1+w)} \tag{2.14}$$

and

$$-V = w\rho_{w0}a^{-3(1+w)}, \tag{2.15}$$

where ρ_{dm0} and ρ_{w0} are the current energy-density of dark matter and that of a fluid, respectively. Here, the current value of the scale factor is chosen as $a_0 = 1$. Thus,

$$\lambda = -\frac{1}{2}\left\{\rho_{dm0}a^{-3} + (1+w)\rho_{w0}a^{-3(1+w)}\right\}, \tag{2.16}$$

$$V = -w\rho_{w0}a^{-3(1+w)}. \tag{2.17}$$

The specific form of the potential is very complicated even if it can be obtained by solving Eq. (2.5). However, one can easily obtain some approximated forms of $V(\phi)$.

When $w > -1$, an approximated solution of wCDM model is given by

$$a(t) = \left[\frac{3}{4}\kappa^2(\rho_{b0}+\rho_{dm0})\right]^{\frac{1}{3}} t^{\frac{2}{3}} \left\{1 + \left[(1+w)^2\frac{\rho_{w0}}{\rho_{b0}+\rho_{dm0}}\left(\frac{3}{4}\kappa^2(\rho_{b0}+\rho_{dm0})t^2\right)^{-w}\right]^{\frac{1}{3(1+w)}}\right\}, \tag{2.18}$$

where ρ_{b0} is the current energy-density of the baryonic matter, because the solutions of Eq. (2.5) in the matter dominant universe and in the fluid matter dominant universe are given by $a = (3\kappa^2(\rho_{b0}+\rho_{dm0})/4)^{1/3}t^{2/3}$ and $a = [3(1+w)^2\kappa^2\rho_{w0}/4]^{1/(3(1+w))}t^{2/(3(1+w))}$, respectively. Therefore, an approximated form of potential $V(\phi)$ is expressed as

$$V(\phi) = \frac{4}{9\kappa^2\phi^2}\frac{1}{1+\left[(1+w)^2\frac{\rho_{w0}}{\rho_{b0}+\rho_{dm0}}\left(\frac{3}{4}\kappa^2(\rho_{b0}+\rho_{dm0})\phi^2\right)^{-w}\right]^{-\frac{1}{3(1+w)}}}$$
$$\times\left\{-\frac{3w+w^2}{(1+w)^2} + \frac{w^2}{(1+w)^2}\frac{1}{1+\left[(1+w)^2\frac{\rho_{w0}}{\rho_{b0}+\rho_{dm0}}\left(\frac{3}{4}\kappa^2(\rho_{b0}+\rho_{dm0})\phi^2\right)^{-w}\right]^{-\frac{1}{3(1+w)}}}\right\}. \tag{2.19}$$

Eq. (2.18) is a very rough approximation as a kind of the interpolation. To improve the accuracy of the approximation, one can use the current value of the Hubble rate H_0 as:

$$a(t) = \left[\frac{3}{4}\kappa^2(\rho_{b0}+\rho_{dm0})t^2\right]^{\frac{1}{3}} + \left[\frac{3}{4}(1+w)^2\kappa^2\rho_{w0}t^2\right]^{\frac{1}{3(1+w)}} + C(H_0t)^{\frac{2+w}{3(1+w)}}, \tag{2.20}$$

where C is a constant tuned to satisfy $\dot{a}/a = H_0$ when $a(t) = 1$. The value of C is evaluated by solving a simultaneous equation $\{a(t_0) = 1,\, H(t_0) = H_0\}$ with respect to t_0 and C. In the case of $w = -0.7$, $t_0 = 116.752\times 10^8$ years and $C = 0.126265$ are obtained. The comparisons between the exact wCDM model and Eq. (2.20) when $w = -0.7$ and $w = -0.9$ are shown in Fig. 1. It seems that Eq. (2.20) is not a bad approximation when w is large, although only three points $t = +0, t_0, +\infty$ are used as the interpolation points.



Figure 1: Comparison of the time dependence of the scale factor between wCDM model and Eq. (2.20). Thick solid line, solid line, dashed line, and dotted line represent Eq. (2.20) with $w = -0.7$, Eq. (2.20) with $w = -0.9$, wCDM model with $w = -0.7$, and wCDM model with $w = -0.9$, respectively. $H_0 = 74$ km/s/Mpc, $\Omega_{m0} = 0.3$, and $\Omega_{w0} = 0.7$ are assumed.

2. *Reconstruction with respect to Om*

One can rewrite Eqs. (2.11) and (2.12) by using $Om(z)$ and $Om'(z)$ instead of $H(t)$ and $H'(t)$ as

$$\lambda = \frac{H_0^2}{\kappa^2}\left\{-3/2(1+z)^3 Om - \frac{1}{2}(1+z)^4 Om' + \frac{1}{2}(1+z)Om'\right\} + \frac{1}{2}\rho_{b0}(1+z)^3, \tag{2.21}$$

$$V = \frac{H_0^2}{\kappa^2}\left\{3(1-Om) - (1+z)^4 Om' + (1+z)Om'\right\}, \tag{2.22}$$

where ρ_{b0} is current energy-density of the baryons. $Om(z)$ is defined by [10, 11]

$$Om(z) = \frac{(H(z)/H_0)^2 - 1}{(1+z)^3 - 1}. \tag{2.23}$$

In the case of the ΛCDM model, $Om(z)$ is in accord with the current matter fraction Ω_{m0}. However, in the recent paper[15], a discrepancy between CMB observations (Planck+WP) and BAO observations is pointed out. It is pointed out that the value of Omh^2 is larger for larger redshift. The explicit values of $\Omega_{m0}h^2$ for Planck+WP and Omh^2 for BAO observations are $\Omega_{m0}h^2 = 0.1426 \pm 0.0025$[16] and $Omh^2 \approx 0.122 \pm 0.01$[15], respectively. Of course, these values are incompatible in the ΛCDM model, because Omh^2 is constant with respect to redshift. However, we can construct a model that satisfies these conditions. If we consider the following scale factor as a solution of Eqs. (2.5) and (2.6):

$$
\begin{aligned}
a(t) &= \left(\frac{g(t)}{1-g(t)}\right)^{1/3} \sinh^{2/3}\left[\frac{3}{2}\sqrt{1-g(t)}H_0 t\right], \\
g(t) &= \left(0.122 + 0.04\frac{t_{CMB}}{t} - 0.02\frac{t_{CMB}^2}{t^2}\right)/h^2,
\end{aligned}
\tag{2.24}
$$

where t_{CMB} is a time of $z = 1089$, then $Omh^2 \approx 0.122$ for small redshift and $Omh^2 = 0.142$ at $z = 1089$ is realized. The potential which causes such an expansion history is given by using Eq. (2.13), however, it is not written down explicitly, being very complicated. Similarly, we can reconstruct any requested universe history. In the next section, the evolution of the matter density perturbations is studied.

III. COSMOLOGICAL PERTURBATIONS WITH ACCOUNT OF MATTER

In this section, we investigate the growth rate of the matter density perturbations (for general review of cosmological perturbations theory, see [12–14]). When we use the metric of the Newtonian gauge, $ds^2 = (-1+2\Phi)dt^2 + (1+2\Psi)a^2(t)\sum_{i=1}^{3} dx^i dx^i$, then (00), $(0i) = (i0)$, (ij), $i \neq j$, and (ii) elements of the linearized Einstein equation in the Fourier space are represented as follows, respectively:

$$
-6H^2\Phi - 2\frac{k^2}{a^2}\Psi - 6H\partial_0\Psi = -\kappa^2\delta\rho + 2\kappa^2\delta\lambda + 2\kappa^2\lambda\Phi + 2\kappa^2\lambda\delta\dot{\phi} - \kappa^2 V_{,\phi}\delta\phi, \tag{3.1}
$$

$$
2(H\Phi + \partial_0\Psi) = \kappa^2(\rho + p)\delta u + 2\kappa^2\lambda\delta\phi, \tag{3.2}
$$

$$
\Phi - \Psi = 0, \tag{3.3}
$$

$$
\begin{aligned}
\left(\frac{k^2}{a^2} + \frac{\partial_i\partial_i}{a^2} - 2H\partial_0 - 4\dot{H} - 6H^2\right)\Phi - \left(\frac{k^2}{a^2} + \frac{\partial_i\partial_i}{a^2} + 2\partial_0\partial_0 + 6H\partial_0\right)\Psi &= \\
\kappa^2 c_{\mathrm{s}}^2\delta\rho - 2\kappa^2\lambda(\Phi + \delta\dot{\phi}) - \kappa^2 V_{,\phi}\delta\phi&,
\end{aligned}
\tag{3.4}
$$

where we defined the sound speed by $c_{\mathrm{s}}^2 \equiv \delta p/\delta\rho$, the matter density perturbation by $\delta \equiv \delta\rho/\rho$ and the scalar perturbation of the four velocity as $\partial_i\delta u$. The linearized equations of (2.3) and (2.4) are given as follows:

$$
\delta\dot{\lambda} + 3H\delta\lambda + \lambda\delta\ddot{\phi} + (\dot{\lambda} + 3H\lambda)\delta\dot{\phi} + \left(\lambda\frac{k^2}{a^2} - \frac{1}{2}V_{,\phi\phi}\right)\delta\phi
$$

$$+\lambda\dot{\Phi} + 2(\dot{\lambda} + 3H\lambda)\Phi + 3\lambda\dot{\Psi} = 0, \tag{3.5}$$

$$\dot{\phi}\delta\dot{\phi} + \dot{\phi}^2\Phi = 0. \tag{3.6}$$

From the perturbation of the equation of continuity, $\nabla_\mu T^{\mu\nu} = 0$, one gets

$$\delta\dot{\rho} + 3H(\delta\rho + \delta p) + a^{-2}\partial_i\{(\rho + p)\partial_i\delta u\} + 3\dot{\Psi}(\rho + p) = 0\,, \tag{3.7}$$

$$a^{-3}\partial_0\{a^3(\rho + p)\partial_i\delta u\} + c_{\mathrm{s}}^2\partial_i\delta\rho - (\rho + p)\partial_i\Phi = 0\,. \tag{3.8}$$

In the following, we treat the equation of state parameter and the sound speed of the matter as $w = c_{\mathrm{s}} = 0$ by assuming the era after matter dominance.

A. Mimetic dark matter

First, we consider the case $V'(\phi) = 0$ as a simple example. Then, the field λ behaves as dark matter in the flat FLRW space-time[5]. Let us investigate whether the field λ also behaves as dark matter when we consider cosmological perturbations. First, Eqs. (3.6) and (3.3) give the relation $\Phi = \Psi = -\delta\dot{\phi}$. By applying the subhorizon approximation, $\partial_0 \sim H$ and $H^2 \ll k^2/a^2$, Eqs. (3.1) and (3.5), then, yield the following equations:

$$2\frac{k^2}{a^2}\delta\dot{\phi} \simeq -\kappa^2\delta\rho + 2\kappa^2\delta\lambda, \tag{3.9}$$

$$\delta\dot{\lambda} + 3H\delta\lambda + \lambda\frac{k^2}{a^2}\delta\phi \simeq 0, \tag{3.10}$$

Differentiating Eq. (3.10) with respect to t gives

$$\ddot{\delta}_\lambda + 2H\dot{\delta}_\lambda + \frac{k^2}{a^2}\delta\dot{\phi} \simeq 0, \tag{3.11}$$

where $\delta_\lambda \equiv \delta\lambda/\lambda$. In the case of $w = c_s^2 = 0$, we can obtain the following equation by using Eqs. (3.7), (3.8),

$$\ddot{\delta} + 2H\dot{\delta} + \frac{k^2}{a^2}\delta\dot{\phi} \simeq 0. \tag{3.12}$$

Taking into account $\rho, \lambda \propto a^{-3}$, we obtain

$$\ddot{\delta}_{\mathrm{tot}} + 2H\dot{\delta}_{\mathrm{tot}} + \frac{k^2}{a^2}\delta\dot{\phi} \simeq 0, \tag{3.13}$$

where $\rho_{\mathrm{tot}} \equiv \rho - 2\lambda$ and $\delta_{\mathrm{tot}} \equiv (\delta\rho - 2\delta\lambda)/(\rho - 2\lambda)$. By using Eq. (3.9), we finally obtain

$$\ddot{\delta}_{\mathrm{tot}} + 2H\dot{\delta}_{\mathrm{tot}} - \frac{\kappa^2}{2}\rho_{\mathrm{tot}}\delta_{\mathrm{tot}} \simeq 0. \tag{3.14}$$

Now, one can recognize the equivalence between mimetic dark matter and fluid dark matter at the linear perturbation level because Eq. (3.14) is same as that of the ΛCDM model. The equivalence of mimetic dark matter and fluid dark matter had been shown in Ref. [7].

B. Mimetic matter with potential

In this section, we consider more involved case $V_{,\phi} \neq 0$. The perturbation equations (3.2), (3.4), and (3.7) can be simplified as follows by using Eqs. (3.4) and (3.6):

$$-2\delta\ddot{\phi} - 2H\delta\dot{\phi} = \kappa^2\rho\delta u + 2\kappa^2\lambda\delta\phi, \tag{3.15}$$

$$2\delta\dddot{\phi} + 8H\delta\ddot{\phi} + (4\dot{H} + 6H^2)\delta\dot{\phi} = -\kappa^2 V_{,\phi}\delta\phi, \tag{3.16}$$

$$\delta\dot{\rho} + 3H\delta\rho - \frac{k^2}{a^2}\rho\delta u - 3\rho\delta\ddot{\phi} = 0\,. \tag{3.17}$$

Eliminating the term proportional to δu from Eqs. (3.15) and (3.17) gives

$$\dot{\delta} + \left(\frac{2}{\kappa^2\rho}\frac{k^2}{a^2} - 3\right)\delta\ddot{\phi} + \frac{2H}{\kappa^2\rho}\frac{k^2}{a^2}\delta\dot{\phi} + \frac{2\lambda}{\rho}\frac{k^2}{a^2}\delta\phi = 0. \tag{3.18}$$

Then, we can derive the following evolution equation of the matter density perturbation:

$$\begin{aligned}\ddddot{\delta} &+ \left(7H - \frac{\dot{\lambda}}{\lambda} + O\left(\frac{a^2H^2}{k^2}\right)\right)\dddot{\delta} + \left(16H^2 - 4\kappa^2(\rho - 2\lambda) - 5H\frac{\dot{\lambda}}{\lambda} + O\left(\frac{a^2H^2}{k^2}\right)\right)\ddot{\delta} \\ &+ \frac{3}{2}\left(8H^3 - 3\kappa^2H(\rho - 4\lambda) - 4H^2\frac{\dot{\lambda}}{\lambda} + \kappa^2\rho\frac{\dot{\lambda}}{\lambda} + O\left(\frac{a^2H^2}{k^2}\right)\right)\dot{\delta} = 0.\end{aligned} \tag{3.19}$$

Eq. (3.19) is quite different from that in quintessence model. There are no oscillating solutions, moreover, the quasi-static solutions are modified. In fact, if we assume

$$\ddot{\delta} + 2H\dot{\delta} - \frac{\kappa^2}{2}\rho\delta = 0, \tag{3.20}$$

then

$$\begin{aligned}\ddddot{\delta} &+ \left(7H - \frac{\dot{\lambda}}{\lambda}\right)\dddot{\delta} + \left(16H^2 - 4\kappa^2(\rho - 2\lambda) - 5H\frac{\dot{\lambda}}{\lambda}\right)\ddot{\delta} \\ &+ \frac{3}{2}\left(8H^3 - 3\kappa^2H(\rho - 4\lambda) - 4H^2\frac{\dot{\lambda}}{\lambda} + \kappa^2\rho\frac{\dot{\lambda}}{\lambda}\right)\dot{\delta} - \frac{\kappa^4}{2}\rho\lambda\delta = 0.\end{aligned} \tag{3.21}$$

Therefore, the solution which satisfies Eq. (3.20) is not the solution of Eq. (3.19) except for the cases $\rho = 0$ or $\lambda = 0$ or $\delta = 0$. It is necessary to be careful because there is an ambiguity to express Eq. (3.21) due to adding the terms proportional to Eq. (3.20) or the derivatives of Eq. (3.20) are allowed. In k-essence model, the growth equation of the matter density perturbation is also represented by fourth order equation[17] as in Eq. (3.19). However, in general, the sound speed read off from the field equation (3.5) is not zero in the k-essence model, so there are no modifications to the quasi-static solution. Thus, the modification to the quasi-static solution in Eq. (3.19) is the peculiar feature in this theory. However, in the matter dominant era or in λ dominant era, the term $-\kappa^4\rho\lambda\delta/2$ is less than the other terms in Eq. (3.21). Therefore, we need to consider the era where there is almost same amount of dark energy and dark matter such as the current universe to distinguish Eq. (3.19) from Eq. (3.21). Whereas, in the case of $V_{,\phi} = 0$, Eq. (3.19) recovers the differentiated equation of (3.14), correctly.

The procedure to obtain Eq. (3.19) is

1. differentiating Eq. (3.18) with respect to t
2. eliminating the terms proportional to $\delta\dddot{\phi}$ from the equation derived at the first step by using Eq. (3.16)
3. differentiating the equation derived at the second step with respect to t
4. eliminating the terms proportional to $\delta\dddot{\phi}$ from the equation derived at the third step by using Eq. (3.16)
5. differentiating the equation derived at the fourth step with respect to t
6. eliminating the terms proportional to $\delta\dddot{\phi}$ from the equation derived at the fifth step by using Eq. (3.16)
7. eliminating the terms proportional to $\delta\ddot{\phi}$ from the equations derived at the second, fourth, and sixth step by using Eq. (3.18)
8. eliminating the terms proportional to $\delta\dot{\phi}$ and $\delta\phi$ by using the equations derived at the seventh step.



Figure 2: Redshift dependence of the growth rate function $f(z)$ in mimetic wCDM. The dot-dash line, the solid line, and the dashed line express $f(z)$ in the cases $w = -1.2$, $w = -1$, and $w = -0.8$, respectively. $\Omega_{w0} = 0.73$, $\Omega_{dm0} = 0.23$, and $\Omega_{b0} = 0.04$ are assumed to depict the above figure. Initial conditions are $f''(100) = f'(100) = 0$ and $f(100) = 1$.

1. Mimetic wCDM

Let us evaluate the difference between Eqs. (3.19) and (3.20) in the case that mimetic matter behaves like wCDM model. For a correct evaluation, we will not use Eq. (2.19) but use Eqs. (2.16) and (2.17). If we utilize $N \equiv \ln a$ instead of time t, we can describe all the functions in Eq. (3.19) as the functions of a by using Eqs. (2.5), (2.6), (2.10), (2.16), and (2.17). Therefore, Eq. (3.19) can be solved at least by numerical calculations. Fig. 2 shows the changes of the growth rate function $f(z) \equiv d \ln \delta(z)/dN$, with respect to z, where $z = -1 + 1/a = -1 + \mathrm{e}^{-N}$. Figs. 3 and 4 express the differences between mimetic wCDM and wCDM model in the growth rate function. Here, $\Omega_{w0} = 0.73$, $\Omega_{dm0} = 0.23$, and $\Omega_{b0} = 0.04$, where $\Omega_{X0} \equiv \kappa^2 \rho_{X0}/(3H_0^2)$ is the current energy fraction of the matter X, are assumed to depict the figures. As the initial conditions, $f''(100) = f'(100) = 0$ and $f(100) = 1$ are used, because, in the matter dominant era, i.e. $V \simeq const.$, we could use Eq. (3.14) and can obtain $f''(z) = f'(z) = 0$ and $f(z) = 1$. As noted in Sec. III A and at the beginning of Sec. III B, there is no difference between mimetic model and wCDM model when $w = -1$. However,



Figure 3: Difference between mimetic wCDM and wCDM model in the growth rate function $f(z)$ when $w = -0.8$.

in the case $w \neq -1$ & $w < 0$, there is difference which come from the term $-\kappa^4 \rho \lambda \delta / 2$ in Eq. (3.21). One can see in Figs. 3 and 4 that the differences are small but surely exist. The term $-\kappa^4 \rho \lambda \delta / 2$ is not relevant at the matter dominant era, so we can also find that the differences become small as redshift z becomes large.

2. Dynamical Om(z)

The behavior of the growth rate function $f(z)$ when the background solution of the Universe is given by Eq. (2.24) is expressed in Fig. 5. $H_0 = 100 h kms^{-1} Mpc^{-1}$, $h = 0.7$ is assumed to depict Fig. 5. Although $Om(z)$ is a smoothly changing function from $0.142/h^2$ to $0.122/h^2$ when we consider $z < 1089$, however, the growth rate function $f(z)$ is larger than that of the ΛCDM model with $\Omega_{m0} h^2 = 0.142$. This effect comes from the difference between Eqs. (3.19) and (3.20). In fact, if we evaluate $f(z)$ for a constant Om model with $Omh^2 = 0.142$ in Eq. (3.19), we have a larger value than that of the dynamical $Om(z)$ (2.24).



Figure 4: Difference between mimetic wCDM and wCDM model in the growth rate function $f(z)$ when $w = -1.2$.



Figure 5: The growth rate function $f(z)$ when the evolution of the background space-time is described by Eq. (2.24). Solid, Dashed, and Dot-dashed curves represent $f(z)$ in this model, $f(z)$ in the ΛCDM model with $\Omega_{m0}h^2 = 0.142$, and $f(z)$ in the ΛCDM model with $\Omega_{m0}h^2 = 0.122$, respectively. Initial conditions are $f''(100) = f'(100) = 0$ and $f(100) = 1$.

IV. ON THE POSSIBILITY OF THE PHANTOM – NON-PHANTOM TRANSITION

A. Reconstruction of the background evolution

In the quintessence model, the value of EoS parameter $w = p_\phi/\rho_\phi$ is constrained as $-1 < w < 1$ so that phantom behavior $w < -1$ cannot be realized. Here, we formally consider the possibility to construct a model, which can describe the transition from $-1 < w$ to $w < -1$ or from $w < -1$ to $-1 < w$. If the Hubble rate $H(t)$ is given by the following function,

$$H(t) = \frac{\alpha}{3}(T_0 + t)^3 - \beta(T_0 + t) + \gamma,\ \gamma = -\frac{\alpha}{3}T_0^3 + \beta T_0, \tag{4.1}$$

$$H(t) = h_0 + h_1 \sin(\nu t), \tag{4.2}$$

$$H(t) = c_0 \left(\frac{1}{t} + \frac{1}{t_s - t}\right), \tag{4.3}$$

or

$$H(t) = t_2 \left(\frac{1}{t_0^2 - t^2} + \frac{1}{t_1^2 + t^2}\right), \tag{4.4}$$

then for the case, $0 < t < t_-$ or $t_+ < t$, $h_1\nu\cos(\nu t) > 0$, $t_s/2 < t < t_s$, or $t'_- < t < 0$ or $t > t'_+$, where $t_\pm \equiv -T_0 \pm \sqrt{\beta/\alpha}$ and $t'_\pm \equiv \pm\sqrt{(t_0^2 - t_1^2)/2}$, effective EoS parameter $w_{\text{eff}} = -2\dot{H}/(3H^2) - 1$ satisfies $w_{\text{eff}} < -1$, respectively[18]. Here, T_0, h_1, and ν are real valued constants and $\alpha, \beta, h_0, c_0, t_s, t_0, t_1 < t_0$ and t_2 are positive constants. Eq. (4.1) could also describe the unified model of the inflation and dark energy because there are two regimes $0 < t < t_-$ and $t_+ < t$ satisfying $w_{\text{eff}} < -1$, the era $0 < t < t_-$ and the era $t_+ < t$ correspond to the inflation and a dominance of dark energy, respectively. The forms of the potential which cause the above expansion rates of the Universe are obtained by using Eq. (2.13) as

$$\begin{aligned} V(\phi) = \frac{1}{\kappa^2}\Bigg[&\frac{\alpha^2}{3}(T_0 + \phi)^6 - 2\alpha\beta(T_0 + \phi)^4 + 2\alpha\gamma(T_0 + \phi)^3 \\ &+(2\alpha + 3\beta^2)(T_0 + \phi)^2 - 6\beta\gamma(T_0 + \phi) + 3\gamma^2 - 2\beta\Bigg], \end{aligned} \tag{4.5}$$

$$V(\phi) = \frac{1}{\kappa^2}\left[3h_0^2 + 6h_0 h_1 \sin(\nu\phi) + 2h_1\nu\cos(\nu\phi) + 3h_1^2\sin^2(\nu\phi)\right], \tag{4.6}$$

$$V(\phi) = \frac{1}{\kappa^2}\left[\frac{3c_0^2 - 2c_0}{\phi^2} + \frac{6c_0^2}{\phi(t_s - \phi)} + \frac{2c_0 + 3c_0^2}{(t_s - \phi)^2}\right], \tag{4.7}$$

and

$$V(\phi) = \frac{1}{\kappa^2}\left[\frac{4t_2\phi + 3t_2^2}{(t_0^2 - \phi^2)} + \frac{6t_2^2}{(t_0^2 - \phi^2)(t_1^2 + \phi^2)} + \frac{3t_2^2 - 4t_2\phi}{(t_1^2 + \phi^2)^2}\right], \tag{4.8}$$

respectively. Here, EoS parameter of the matter is assumed to be 0. As seen in Eq. (2.16), the phantom crossings in these examples are realized by the transition from $\lambda > 0$ to $\lambda < 0$ when there is no usual matter. Hence, in mimetic matter model with potential the phantom-divide transitions maybe realized by proper choice of the scalar potential.

B. Growth of the matter density contrast in phantom crossing model

Let us consider the evolution behavior of the matter density contrast in the model with Eq. (4.7) as an example. In this model, the Universe will experience the singular evolution in $H(t)$ if t_s is larger than the current time. It is known that there may occur several finite-time future singularity scenarios of the Universe as [19–21] which are classified as [20]:

- Type I (”Big Rip”) : For $t \to t_s$, $a \to \infty$, $\rho_{\mathrm{eff}} \to \infty$, and $|p_{\mathrm{eff}}| \to \infty$. This also includes the case of ρ_{eff}, p_{eff} being finite at t_s.
- Type II (”Sudden”) : For $t \to t_s$, $a \to a_s$, $\rho_{\mathrm{eff}} \to \rho_s$, and $|p_{\mathrm{eff}}| \to \infty$.
- Type III : For $t \to t_s$, $a \to a_s$, $\rho_{\mathrm{eff}} \to \infty$, and $|p_{\mathrm{eff}}| \to \infty$.
- Type IV : For $t \to t_s$, $a \to a_s$, $\rho_{\mathrm{eff}} \to 0$, $|p_{\mathrm{eff}}| \to 0$, and higher derivatives of H diverge. This also includes the case in which p_{eff} (ρ_{eff}) or both of p_{eff} and ρ_{eff} tend to some finite values, while higher derivatives of H diverge.

Singularity in the case under consideration corresponds to Type I or Type III. When $c_0 = 2/3$ and $t_s = 4 \times 10^{10}$ years, we have a solution of Eq. (4.3) as

$$a(t) = c_1 \left(\frac{t}{t_s - t} \right)^{\frac{2}{3}} , \tag{4.9}$$

where c_1 is an integration constant. It is chosen as $c_1 = 1.6$ to comply with the observed background evolution of the Universe. In this case, w_{eff} is simply expressed as $w_{\mathrm{eff}} = -2t/t_s$. The evolution behavior of the matter density perturbation is changed around the singularity because the approximation $|a^2H^2/k^2| \ll 1$ no longer holds in Eq. (3.19). In Fig. 6, the behavior of the growth rate function $f(z) = d \ln \delta(z)/dN(z)$ until the current time is expressed. On the other hand, the future evolution of $f(t)$ is expressed in Fig. 7. Fig. 7 shows singular behavior in $f(t)$ around $t = 2.7 \times 10^{10}$ years, although the singularity in $a(t)$ and $H(t)$ occurs at $t = 4\times10^{10}$ years. The reason why such a behavior of $f(t)$ occurs is instability



Figure 6: The growth rate function $f(z)$ in the model (4.7)

of the perturbations of phantom matter even at the classical level. To be concrete, the flip of the sign of the kinetic term would cause the instability. A decay of perturbations much earlier of future singularity found in [32] is just the same as this instability (for discussion of problems of phantom-divide transitions in perturbations, see [22–24]).

V. CONCLUSIONS

In this paper, we have investigated the evolution of the matter density perturbation in mimetic matter model. In Sec. II, the reconstruction, the way to construct a model which can describe arbitrary evolution history of the Universe, has been explicitly shown and some examples, including mimetic wCDM model and evolving Om(z) model, have been given. At the beginning of Sec. III, the growth rate of the matter density perturbation has been investigated when potential term is not dynamical. As a result, it has been shown that mimetic dark matter also behaves as CDM at the perturbation level. In the case of dynamical potential, it has been shown that the differential equation of the matter density perturbation is changed. In general, this equation is different from that in the ΛCDM model and the quasi-static equation of the matter density perturbation in quintessence model. We have presented the behavior of the growth rate function in mimetic wCDM model and in



Figure 7: The future evolution of the growth rate function $f(t)$ in the model (4.7)

a model of dynamical $Om(z)$ as examples. The difference seems not to happen in mimetic wCDM, but it clearly appears in dynamical $Om(z)$. In Sec. IV, we have proved the possibility to realize phantom – non-phantom transition in mimetic matter model. It has been shown that phantom – non-phantom transition is possible because the sign of kinetic term of the scalar field is determined by the sign of the Lagrange multiplier. However, usually phantom brings an instability. Such an instability occurs not only as quantum effect but also as classical effect[25–31], and it breaks the perturbation theory as is explicitly demonstrated.

It would be interesting to study the accelerating universe evolution and cosmological perturbations in the generalizations of mimetic matter model which include mimetic $F(R)$ gravity [33, 36], higher-derivative mimetic model[34] or its further generalizations[35]. In this case, we have not only scalar potential but also extra function like $F(R)$ or higher-derivative term. Hence, the cosmological reconstruction can be executed quite easily. This will be studied elsewhere.

Acknowledgments

JM would like to acknowledge a suggestion from A. A. Starobinsky. The work was supported by the Russian Government Program of Competitive Growth of Kazan Federal University and, partially, by the Russian Foundation for Basic Research grants No. 14-02-00598. The research by SDO has been supported in part by MINECO (Spain), projects FIS2010-15640 and FIS2013-44881.

---

[1] K. Bamba, S. Capozziello, S. Nojiri and S. D. Odintsov, Astrophys. Space Sci. **342**, 155 (2012) [arXiv:1205.3421 [gr-qc]].

[2] A. Joyce, B. Jain, J. Khoury and M. Trodden, arXiv:1407.0059 [astro-ph.CO]; Y. F. Cai, E. N. Saridakis, M. R. Setare and J. Q. Xia, Phys. Rept. **493** (2010) 1 [arXiv:0909.2776 [hep-th]];

[3] G. Bertone, D. Hooper and J. Silk, Phys. Rept. **405**, 279 (2005) [hep-ph/0404175].

[4] J. L. Feng, Ann. Rev. Astron. Astrophys. **48**, 495 (2010) [arXiv:1003.0904 [astro-ph.CO]].

[5] A. H. Chamseddine and V. Mukhanov, JHEP **1311**, 135 (2013) [arXiv:1308.5410 [astro-ph.CO]].

[6] A. H. Chamseddine, V. Mukhanov and A. Vikman, JCAP **1406**, 017 (2014) [arXiv:1403.3961 [astro-ph.CO]].

[7] E. A. Lim, I. Sawicki, and A. Vikman, JCAP **1005**, 012 (2010) [arXiv:1003.5751 [astro-ph.CO]]; S. Capozziello, J. Matsumoto, S. Nojiri and S. D. Odintsov, Phys. Lett. B **693** (2010) 198 [arXiv:1004.3691 [hep-th]].

[8] A. Golovnev, Phys. Lett. B **728**, 39 (2014) [arXiv:1310.2790 [gr-qc]].

[9] A. O. Barvinsky, JCAP **1401**, no. 01, 014 (2014) [arXiv:1311.3111 [hep-th]].

[10] V. Sahni, A. Shafieloo and A. A. Starobinsky, Phys. Rev. D **78**, 103502 (2008) [arXiv:0807.3548 [astro-ph]].

[11] C. Zunckel and C. Clarkson, Phys. Rev. Lett. **101**, 181301 (2008) [arXiv:0807.4304 [astro-ph]].

[12] V. Mukhanov, "Physical foundations of cosmology," Cambridge, UK: Univ. Pr. (2005) 421 p; V. F. Mukhanov, H. A. Feldman and R. H. Brandenberger, Phys. Rept. **215** (1992) 203.

[13] D. S. Gorbunov and V. A. Rubakov, "Introduction to the theory of the early universe: Cos-

mological perturbations and inflationary theory," Hackensack, USA: World Scientific (2011) 489 p

[14] H. Kodama and M. Sasaki, Prog. Theor. Phys. Suppl. **78** (1984) 1.

[15] V. Sahni, A. Shafieloo and A. A. Starobinsky, Astrophys. J. **793**, no. 2, L40 (2014) [arXiv:1406.2209 [astro-ph.CO]].

[16] P. A. R. Ade *et al.* [Planck Collaboration], Astron. Astrophys. **571**, A16 (2014) [arXiv:1303.5076 [astro-ph.CO]].

[17] K. Bamba, J. Matsumoto and S. Nojiri, Phys. Rev. D **85**, 084026 (2012) [arXiv:1109.1308 [hep-th]].

[18] S. Nojiri and S. D. Odintsov, Gen. Rel. Grav. **38**, 1285 (2006) [hep-th/0506212].

[19] A. A. Starobinsky, Grav. Cosmol. **6**, 157 (2000) [astro-ph/9912054].

[20] S. Nojiri, S. D. Odintsov and S. Tsujikawa, Phys. Rev. D **71**, 063004 (2005) [arXiv:hep-th/0501025].

[21] R. R. Caldwell, M. Kamionkowski and N. N. Weinberg, Phys. Rev. Lett. **91** (2003) 071301 [astro-ph/0302506];J. D. Barrow, Class. Quant. Grav. **21** (2004) L79 [gr-qc/0403084];Class. Quant. Grav. **21** (2004) 5619 [gr-qc/0409062]; V. Gorini, A. Y. Kamenshchik, U. Moschella and V. Pasquier, Phys. Rev. D **69** (2004) 123512 [hep-th/0311111]; S. Nojiri and S. D. Odintsov, Phys. Lett. B **562** (2003) 147 [hep-th/0303117];Phys. Rev. D **72** (2005) 023003 [hep-th/0505215]; P. Singh, M. Sami and N. Dadhich, Phys. Rev. D **68** (2003) 023522 [hep-th/0305110]; V. Faraoni, Phys. Rev. D **68** (2003) 063508 [gr-qc/0307086]; M. P. Dabrowski, T. Stachowiak and M. Szydlowski, Phys. Rev. D **68** (2003) 103519 [hep-th/0307128]; M. Bouhmadi-Lopez, P. F. Gonzalez-Diaz and P. Martin-Moruno, Phys. Lett. B **659** (2008) 1 [gr-qc/0612135]; L. Fernandez-Jambrina and R. Lazkoz, Phys. Rev. D **74** (2006) 064030 [gr-qc/0607073]; K. Bamba, S. Nojiri and S. D. Odintsov, JCAP **10**, 045 (2008) [arXiv:0807.2575 [hep-th]]; H. Ghodsi, M. A. Hendry, M. P. Dabrowski and T. Denkiewicz, Mon. Not. Roy. Astron. Soc. **414** (2011) 2, 1517 [arXiv:1101.3984 [astro-ph.CO]].

[22] A. Vikman, Phys. Rev. D **71**, 023515 (2005) [astro-ph/0407107].

[23] W. Hu, Phys. Rev. D **71**, 047301 (2005) [astro-ph/0410680].

[24] R. R. Caldwell and M. Doran, Phys. Rev. D **72**, 043527 (2005) [astro-ph/0501104].

[25] S. Dubovsky, T. Gregoire, A. Nicolis and R. Rattazzi, JHEP **0603**, 025 (2006) [hep-th/0512260].

[26] R. P. Woodard, Lect. Notes Phys. **720**, 403 (2007) [astro-ph/0601672].

[27] I. Sawicki and A. Vikman, Phys. Rev. D **87**, no. 6, 067301 (2013) [arXiv:1209.2961 [astro-ph.CO]].

[28] J. M. Cline, S. Jeon and G. D. Moore, Phys. Rev. D **70**, 043543 (2004) [hep-ph/0311312].

[29] J. Garriga and A. Vilenkin, JCAP **1301**, 036 (2013) [arXiv:1202.1239 [hep-th]].

[30] E. Elizalde, S. Nojiri and S. D. Odintsov, Phys. Rev. D **70**, 043539 (2004) [hep-th/0405034].

[31] S. Nojiri and S. D. Odintsov, Phys. Rev. D **72**, 023003 (2005) [hep-th/0505215].

[32] A. V. Astashenok and S. D. Odintsov, Phys. Lett. B **718**, 1194 (2013) [arXiv:1211.1888 [gr-qc]].

[33] S. Nojiri and S. D. Odintsov, arXiv:1408.3561 [hep-th].

[34] L. Mirzagholi and A. Vikman, arXiv:1412.7136 [gr-qc].

[35] D. Momeni, A. Altaibayeva and R. Myrzakulov, arXiv:1407.5662 [gr-qc].

[36] G. Leon and E. N. Saridakis, arXiv:1501.00488 [gr-qc].